

# SUBJECT INDEX TO VOLUME 40

	Page	Page
<i>a</i> = abstract		
<b>Acetonitrile</b> , effects on charge transport in [Os(bipy) <sub>2</sub> (PVP) <sub>10</sub> Cl]Cl films, <i>a</i>	190	
<b>Acetophenone</b> , derivatives, asymmetric transfer hydrogenation, via Ru(II) complex, <i>a</i>	94	
<b>Acetylcholine</b> , detector	69	
<b>Acetylenes</b> , disubstituted, from C <sub>2</sub> H <sub>2</sub> +aryl iodides, <i>a</i>	141	
<b>Acrylic Acid</b> , hydrogenation, on Pd/PA-Nd <sup>3+</sup> , <i>a</i>	192	
<b>Alcohols</b> , allenic, synthesis, <i>a</i>	141	
allyloxycarbonyl-protected, Pd-catalysed cleavage, <i>a</i>	141	
direct formation from alkenes, <i>a</i>	141	
ethyl, CO <sub>2</sub> hydrogenation, over Rh/SiO <sub>2</sub> +Li salts, <i>a</i>	193	
synthesis from CO <sub>2</sub> -H <sub>2</sub> , on [Rh <sub>10</sub> Se]/TiO <sub>2</sub> clusters, <i>a</i>	92	
from hydrocarbonylation of alkenes, <i>a</i>	141	
hexanol, photoproduction, <i>a</i>	190	
methyl, effects on charge transport in [Os(bipy) <sub>2</sub> (PVP) <sub>10</sub> Cl]Cl films, <i>a</i>	190	
electro-oxidation, at Pt-Ru-Sn, <i>a</i>	137	
on Pt/graphite, <i>a</i>	89	
production from CO <sub>2</sub> via Ru cluster catalysts, <i>a</i>	94	
optically active ( <i>E</i> )-allylic, ( <i>E,E</i> )-dienylic and allenic alcohols, synthesis, <i>a</i>	141	
( <i>S</i> )-enriched, > 90% yield, < 92% ee, production, <i>a</i>	94	
unsaturated, from cinnamaldehyde hydrogenation, <i>a</i>	192	
hydroformylation, by Rh(acac)(CO) <sub>2</sub> +phosphine ligands, <i>a</i>	141	
<b><math>\alpha</math>-Aldehydes</b> , from methylacrylate, via water soluble Rh phosphine PNS/PC, <i>a</i>	93	
<b>Aldehydes</b> , aliphatic, detector, <i>a</i>	42	
fast production from butane, over Pt-Rh single gauze	125	
<b>Alkanes</b> , dehydrogenation, by IrClH <sub>2</sub> (PPr <sub>3</sub> ) <sub>2</sub> , to alkenes, <i>a</i>	141	
hydroxylation, with O <sub>2</sub> and Ru polyoxometalate, <i>a</i>	44	
light, C-C, C-H bond activation, by Pt-Cu/Al <sub>2</sub> O <sub>3</sub> , <i>a</i>	91	
oxidation, by RuF <sub>16</sub> Pc in NaX zeolite catalyst	37	
rapid conversions over Pt-Rh single gauzes	125	
<b>Alkenes</b> , dihydroxylation, <i>a</i>	194	
from alkanes dehydrogenation, by IrClH <sub>2</sub> (PPr <sub>3</sub> ) <sub>2</sub> , <i>a</i>	141	
for hydrocarbonylation to alcohols, <i>a</i>	141	
<b>Alkylation</b> , allylic, by cinnamyl ethyl carbonate, by Pd/TPPTS, <i>a</i>	193	
<b>2-Alkynylbenzaldehyde</b> , carbonylation, <i>a</i>	193	
<b>(<i>E</i>)-Allylic Alcohol</b> , preparation via CpRu(PPh <sub>3</sub> ) <sub>2</sub> Cl, <i>a</i>	194	
<b>Bis(<math>\alpha</math>-Amino Acids)</b> , synthesis, via Pd(PPh <sub>3</sub> ) <sub>4</sub> , <i>a</i>	44	
<b>Ammonia</b> , amperometric detector, <i>a</i>	138	
detection by Pt-doped CdSnO <sub>3</sub> , <i>a</i>	191	
for propionitrile formation, <i>a</i>	92	
synthesis, on Ru clusters/zeolite, <i>a</i>	140	
on Ru/fullerene catalyst, <i>a</i>	43	
on Ru/lanthanide oxide catalysts, <i>a</i>	140	
<b>Ammoxidation</b> , CH <sub>4</sub> , on Pt/Al <sub>2</sub> O <sub>3</sub> , Pt-Rh gauzes, <i>a</i>	192	
<b>Anglo American Platinum Corporation</b>	2	
<b>Aromatisation</b> , propane, on Ga/SiO <sub>2</sub> catalyst in Pd membrane reactor, <i>a</i>	192	
<b>Ascorbic Acid</b> , effect on H <sub>2</sub> O <sub>2</sub> detection, <i>a</i>	191	
<b>Azobenzene</b> , for indole formation, <i>a</i>	193	
<b>Barium</b> , effects on 3-way automotive catalysts, <i>a</i>	92, 192	
<b>Benzocyclobutenes</b> , preparation, with Pd(PPh <sub>3</sub> ) <sub>4</sub> + K <sub>2</sub> CO <sub>3</sub> , <i>a</i>	193	
<b>Biological Systems</b> , PGMs in, book reviews	117, 160	
<b>Book Reviews</b> , "Catalysis by Metals and Alloys"	70	
"Encyclopedia of Chemical Technology"	174	
"Fuel Cells and Their Applications"	159	
"Mixed Ions in Biological Systems, Volume 32: Interactions of Metal Ions with Nucleotides, Nucleic Acid, and Their Constituents"	117	
"Metal Ions in Biological Systems, Volume 33: Probing of Nucleic Acids by Metal Ion Complexes of Small molecules"	160	
"Palladium Reagents and Catalysts-Innovations in Organic Synthesis"	126	
<b>1,3-Butadiene</b> , hydrogenation, over Co-Pd/Al <sub>2</sub> O <sub>3</sub> , <i>a</i>	43	
<b>Butylene</b> , rapid production from butane, over Pt-Rh single gauze	125	
<b>C-H Bonds</b> , in Ru <sub>2</sub> (CO) <sub>8</sub> (C <sub>9</sub> H <sub>10</sub> ), activation via Ru-Ru bond breaking, <i>a</i>	136	
<b>Capacitors</b> , (Ba, Sr)TiO <sub>3</sub> thin film, <i>a</i>	45	
<b>Carbon<sub>60</sub></b> , fullerene pgm complexes	23	
<b>Carbon Oxides</b> , CO <sub>2</sub> hydrogenation, to EtOH, over Rh/SiO <sub>2</sub> +Li salts, <i>a</i>	193	
MeOH, via Ru cluster anions, <i>a</i>	94	
via Rh complex, <i>a</i>	93	
CO <sub>2</sub> + H <sub>2</sub> , for EtOH synthesis, <i>a</i>	92	
CO, CO <sub>2</sub> , reaction kinetics, on polycrystalline Pt electrocatalyst in fuel cell, <i>a</i>	142	
CO-H <sub>2</sub> , electro-oxidation, on (110), (110) surfaces of Pt <sub>2</sub> Sn, structural effects, <i>a</i>	90	
for copolymerisation with ethylene, <i>a</i>	93	
hydrogenation, on Fe+Pt/SiO <sub>2</sub> , <i>a</i>	43	
on Mn-Ru/Al <sub>2</sub> O <sub>3</sub> catalysts, <i>a</i>	140	
on Pt/Co/Al <sub>2</sub> O <sub>3</sub> , <i>a</i>	43	
oxidation, on La <sub>0.8</sub> Sr <sub>0.2</sub> Al <sub>0.8</sub> Cu <sub>0.1</sub> Ru <sub>0.1</sub> O <sub>3</sub> , <i>a</i>	193	
on Pt/CeO <sub>2</sub> , kinetic modelling, <i>a</i>	139	
reactions with Pd-Cu, <i>a</i>	38	
CO-NO reaction, on La <sub>0.8</sub> Sr <sub>0.2</sub> Al <sub>0.8</sub> Cu <sub>0.1</sub> Ru <sub>0.1</sub> O <sub>3</sub> , <i>a</i>	193	
<b>Carbonylation</b> , 2-alkynylbenzaldehyde, via water-gas shift reaction, by Rh <sub>6</sub> (CO) <sub>16</sub> , <i>a</i>	193	
<b>Carboplatin</b> , photostudies, <i>a</i>	138	
<b>Catalysis</b> , aqueous phase, with Rh complex, <i>a</i>	193	
with supported Pd complex, <i>a</i>	193	
conference, 2nd European Congress	19	
heterogeneous, <i>a</i>	42-43, 91-93, 139-140, 191-193	
homogeneous, <i>a</i>	43-44, 93-94, 140-141, 193-194	
conference review	161	
Heck-type cyclisations, <i>a</i>	140	
liquid-phase hydrogenation	8	
Rh supramolecular, enzymatic catalytic reactions, <i>a</i>	93	
<b>Catalysts</b> , see also <b>Photocatalysts</b>		
automotive, 3-way, Pd, CeO <sub>2</sub> effects, <i>a</i>	192	
La, Ba effects, <i>a</i>	92, 192	
Pt-Rh, synergism, <i>a</i>	139	
Pt-Rh/Al <sub>2</sub> O <sub>3</sub> , CeO <sub>2</sub> promoted, <i>a</i>	92	
<b>Catalysts, Iridium</b> , Ir species in H-ZSM-5 zeolite channels, redox, carbonylation chemistry, <i>a</i>	93	
Ir/HY zeolite, Pt/SiO <sub>2</sub> -Al <sub>2</sub> O <sub>3</sub> , for diesel and DBT hydrodesulphurisation, <i>a</i>	139	
<b>Iridium Complexes</b> , for activation of C-H bonds IrClH <sub>2</sub> (PPr <sub>3</sub> ) <sub>2</sub> , for alkane dehydrogenation, <i>a</i>	141	
<b>Osmium</b> , Os, olefin C=C bond fission, in presence of K osmate, and periodic acid, <i>a</i>	90	
<b>Osmium Complexes</b> , OsCl <sub>2</sub> (PPh <sub>3</sub> ) <sub>3</sub> , for cyclopropanation reactions, <i>a</i>	44	
<b>Palladium</b> , Pd colloids, for microcontact printing Pd/ACF, for liquid-phase hydrogenation of olefinic C <sub>6</sub> hydrocarbons, <i>a</i>	140	
Pd/CeO <sub>2</sub> , conversions, <i>a</i>	192	
Pd/ $\gamma$ -Al <sub>2</sub> O <sub>3</sub> , automotive, Ba effects, <i>a</i>	92	
Co-Pd/Al <sub>2</sub> O <sub>3</sub> , 1,3-butadiene hydrogenation, <i>a</i>	43	
Pd metallic/stainless steel flakes, for <i>m</i> -xylene combustion, <i>a</i>	43	
Pd thin film/PbSrZrO, resonance oscillation giving enhanced catalytic activity	25	
Pd/TPPTS, in-situ generated for dynes, indolic and furanic products, Pd(0) intermediate, <i>a</i>	44	
Pd/TPPTS/SiO <sub>2</sub> , water soluble catalyst, for allylic alkylation of cinnamyl ethyl carbonate, <i>a</i>	193	
Pd(OAc) <sub>2</sub> /TPPTS, PdCl <sub>2</sub> /TPPTS, cyclisations, <i>a</i>	140	
<b>Palladium Complexes</b> , (dppp)Pd(OAc) <sub>2</sub> , (dppp)Pd[(C(O)Bu)Cl + <i>tert</i> -butyl alumoxane, for polyketone polymer production, <i>a</i>	93	
Pd[C $\equiv$ C-Si(CH <sub>3</sub> ) <sub>2</sub> (PPh <sub>3</sub> ) <sub>2</sub> ], for ethynyltrimethylsilane polymerisation, <i>a</i>	141	

	Page		Page
<b>Catalysts, Palladium Complexes (contd.)</b>		<b>Catalysts, Rhodium Complexes (contd.)</b>	
homogeneous/support, PdCl <sub>2</sub> -(C <sub>9</sub> H <sub>19</sub> ) <sub>3</sub> N/γ-Al <sub>2</sub> O <sub>3</sub> , for hydrogenations	8	RhCl(PMe <sub>3</sub> ) <sub>3</sub> CO, for hexanal hydrogenation, <i>a</i>	190
Pd(II) system for copolymerisation	78	[RhH(PEt <sub>3</sub> ) <sub>3</sub> ], hydroformylation of hex-1-ene to aldehydes, alcohols, <i>a</i>	141
Pd(OAc) <sub>2</sub> , reaction of bicyclopopylidene and vinyl iodide, cascade of reactions, <i>a</i>	93	Rh(I) supramolecular basket-shape molecule, for dihydroxyarene substituted reactions, <i>a</i>	93
Pd(OAc) <sub>2</sub> -TPPTS, cleavage of allyloxycarbonyl protected alcohols, <i>a</i>	141	Rh(I)(TPPTS)/SiO <sub>2</sub> , for indole production, <i>a</i>	193
[Pd(PPh <sub>3</sub> ) <sub>2</sub> ]Cl <sub>2</sub> +CuI, C <sub>2</sub> H <sub>2</sub> +aryl iodide reaction, <i>a</i>	141	Rh(PPh <sub>3</sub> ) <sub>3</sub> Cl-isocyanide + polymer, alkene, alkyne, nitrile, nitrocompound hydrogenation, <i>a</i>	193
Pd(PPh <sub>3</sub> ) <sub>4</sub> , for bis(α-amino acid) synthesis, <i>a</i>	44	<b>Ruthenium</b> , Ru <sub>3</sub> -C <sub>60</sub> -Cs <sup>+</sup> /SiO <sub>2</sub> , for NH <sub>3</sub> synthesis, <i>a</i>	43
for benzocyclobutenes synthesis, <i>a</i>	193	Ru clusters/zeolite, for NH <sub>3</sub> synthesis, <i>a</i>	140
for polymerisation reactions	78	Ru/La <sub>2</sub> O <sub>3</sub> , for NH <sub>3</sub> synthesis, <i>a</i>	140
(PPh <sub>3</sub> ) <sub>2</sub> PdCl <sub>2</sub> , reaction of 3-( <i>p</i> -nitrobenzoyloxy)- 3-buten-2-one with aryl and vinyl halides, <i>a</i>	193	<b>Ruthenium Alloys</b> , Ru-Me/Al <sub>2</sub> O <sub>3</sub> , Me=Ge, Sn, Pb, for citral, cinnamaldehyde hydrogenation, <i>a</i>	140
<b>Platinum</b> , nanometre Pt/MCM-41, for CO oxidation	35	Ru-Mn/Al <sub>2</sub> O <sub>3</sub> , for CO hydrogenation, <i>a</i>	140
[(PPh <sub>3</sub> )Pt(PPh <sub>3</sub> Au) <sub>6</sub> ](NO <sub>3</sub> ) <sub>2</sub> /SiO <sub>2</sub> , structure, <i>a</i>	139	<b>Ruthenium Complexes</b> ,	
Pt-Bx-Al <sub>2</sub> O <sub>3</sub> , for naphthalene hydrogenation, <i>a</i>	191	[(CO) <sub>2</sub> Rh(μ-Cl)] <sub>2</sub> , for oxabicycloheptenes, cyclopropane ring cleavage, <i>a</i>	43
Pt/Al <sub>2</sub> O <sub>3</sub> , foam monoliths, for HCN synthesis, <i>a</i>	192	Cp*(dppf)RuH, preparation, MV reduction with H <sub>2</sub> , <i>a</i>	94
for NO decomposition + C <sub>2</sub> H <sub>6</sub> , H <sub>2</sub> , <i>a</i>	91	CpRu(PPh <sub>3</sub> ) <sub>2</sub> Cl, for allylic substitution, of allylic cyclic carbonates, <i>a</i>	194
Pt/CeO <sub>2</sub> , kinetic modelling, <i>a</i>	139	Ru <sub>2</sub> (O <sub>2</sub> CCH <sub>3</sub> ) <sub>4</sub> , immobilised on modified supports, for cinnamaldehyde hydrogenation, <i>a</i>	192
Pt/HY zeolite, Pt/SiO <sub>2</sub> -Al <sub>2</sub> O <sub>3</sub> , for diesel, DBT, HDS <i>a</i>	139	Ru <sub>3</sub> (CO) <sub>12</sub> + alkaline iodides, in <i>N</i> -methylpyrrolidone, <i>a</i>	94
Pt/lanthana, for NO, and CO adsorption, <i>a</i>	91	Ru red/Pt, in water electro-oxidation	87
Pt/Na-BEA, for dec-1-ene, dec-5-ene hydrogenation, <i>a</i>	139	RuCl <sub>2</sub> (PPh <sub>3</sub> ) <sub>3</sub> for cyclopropanation reactions, <i>a</i>	44
Pt/TiO <sub>2</sub> , sol-gel, synthesis and characterisation, <i>a</i>	139	RuCl <sub>3</sub> (H <sub>2</sub> O) <sub>3</sub> , for <i>syn</i> -diols, <i>a</i>	194
[Pt(μ-CH <sub>2</sub> COO) <sub>2</sub> ]/SiO <sub>2</sub> , for HCOOH decomposition, <i>a</i>	92	RuCl <sub>3</sub> ·3H <sub>2</sub> O/polycarbonate, NO reduction, water-gas shift reaction, <i>a</i>	93
<b>Platinum Alloys</b> , Pt-Cu/Al <sub>2</sub> O <sub>3</sub> , for alkane C-H, C-C bond activation, <i>a</i>	91	RuF <sub>16</sub> Pc in NaX zeolite catalyst, alkane oxidation	37
Pt-Dy/Al <sub>2</sub> O <sub>3</sub> , for <i>n</i> -hexane, methylcyclopentane conversions, <i>a</i>	192	[RuH <sub>2</sub> (CO)(AsPh <sub>3</sub> ) <sub>3</sub> ], [RuH <sub>2</sub> (CO)(AsPh <sub>3</sub> (L-L))], activity for propionaldehyde hydrogenation, <i>a</i>	194
Pt-Pd/support, S resistance, <i>a</i>	92	RuH <sub>2</sub> (PPh <sub>3</sub> ) <sub>4</sub> , for optically active ( <i>E</i> )-allylic ( <i>E,E</i> )-dienylic, allenic alcohols, synthesis, <i>a</i>	141
Pt-Rh gauzes, for HCN synthesis, <i>a</i>	192	Ru(II), for asymmetric transfer hydrogenation of acetophenone derivatives in propan-2-ol, <i>a</i>	94
Pt-Rh/Al <sub>2</sub> O <sub>3</sub> , CeO <sub>2</sub> -promoted, 3-way, automotive, <i>a</i>	92	[WZnRu <sub>2</sub> III(OH)(H <sub>2</sub> O)(ZnW <sub>6</sub> O <sub>34</sub> ) <sub>2</sub> ] <sup>11-</sup> , for alkane hydroxylation, <i>a</i>	44
Pt-Sn/Al <sub>2</sub> O <sub>3</sub> , sol-gel, preparation, properties, <i>a</i>	192	La <sub>0.8</sub> Sr <sub>0.2</sub> Al <sub>0.8</sub> Cu <sub>0.1</sub> Ru <sub>0.1</sub> O <sub>3</sub> , for CO oxidation, NO-CO reaction, <i>a</i>	193
Pt-Sn/C, for crotonaldehyde hydrogenation, <i>a</i>	139	<b>Catchment Gauzes</b> , Pd alloy, for nitric acid plants	80
<b>Platinum Complexes</b> ,		<b>Cells</b> , Pt/ZrO <sub>2</sub> (Y <sub>2</sub> O <sub>3</sub> )/Pt solid oxide, working electrode function, <i>a</i>	142
PtCl <sub>2</sub> -C <sub>8</sub> H <sub>17</sub> NH <sub>2</sub> -HAl( <i>i</i> -Bu)γ-Al <sub>2</sub> O <sub>3</sub> , for hydrogenations	2	<b>Ceria</b> , promoter, <i>a</i>	92
PtCl <sub>2</sub> , synthesis of 3-oxabicyclo[4.1.0]hept-4-enes, <i>a</i>	43	structural effects on O migration on Rh catalysts, <i>a</i>	92
[PtH(H <sub>2</sub> O)(PMe <sub>3</sub> ) <sub>2</sub> ][OH], acetonitrile hydration chiral, Pt, <i>a</i>	43, 102	<b>Chirality</b> , in Pt, Ru, Os complexes	102
colloidal, for H <sub>2</sub> , photochemical production, <i>a</i>	190	<b>Chlorine</b> , production, on coated Ru-Pt oxide electrodes, <i>a</i>	90
Fe+Pt/SiO <sub>2</sub> , CO + H <sub>2</sub> reactions, <i>a</i>	43	<b>Choline</b> , detector	69
Pt/AAPx, for naphthalene hydrogenation, <i>a</i>	42	<b>Chromium</b> , electrodeposition, via Pt/Ti anodes, <i>a</i>	42
Pt/Al borate, Pt/borate-Al <sub>2</sub> O <sub>3</sub> , resistance to S, <i>a</i>	43	<b>Cinnamaldehyde</b> , hydrogenation, <i>a</i>	140
Pt/Co/Al <sub>2</sub> O <sub>3</sub> , for CO hydrogenation, <i>a</i>	43	over Rh, Ru, Pd complexes, <i>a</i>	192
PtCl <sub>2</sub> (atropisomeric diphosphine)/SnCl <sub>2</sub> , for styrene asymmetric hydroformylation, <i>a</i>	43	<b>Cinnamyl Ethyl Carbonate</b> , allylic alkylation, catalysed by Pd/TPPS/SiO <sub>2</sub> , <i>a</i>	193
<b>Rhodium</b> , liquid-phase hydroformylation catalysts, <i>a</i>	140	<b>Cisplatin</b> , adducts in DNA, crystal structure, <i>a</i>	165
[Rh <sub>10</sub> Se]/TiO <sub>2</sub> clusters, for EtOH synthesis from CO <sub>2</sub> /H <sub>2</sub> , <i>a</i>	92	anti-cancer drug	140
Rh/CeO <sub>2</sub> , ceria structure effect on O migration, <i>a</i>	92	<b>Citral</b> , hydrogenation, <i>a</i>	140
Rh/lanthana, for NO, and CO adsorption, <i>a</i>	91	<b>Clusters</b> , Pt <sub>399</sub> , with ligands, metallic behaviour, <i>a</i>	136
Rh/SiO <sub>2</sub> +Li salts, CO <sub>2</sub> hydrogenation to EtOH, <i>a</i>	193	[(PPh <sub>3</sub> )Pt(PPh <sub>3</sub> Au) <sub>6</sub> ](NO <sub>3</sub> ) <sub>2</sub> , catalyst for H <sub>2</sub> -D <sub>2</sub> equilibration, <i>a</i>	139
Rh/SiO <sub>2</sub> , highly dispersed, from organic Rh precursors via OMCVD, <i>a</i>	92	Ru <sub>3</sub> (CO) <sub>8</sub> (C <sub>6</sub> H <sub>6</sub> ), C-H bond activation on Ru-Ru bond fission, <i>a</i>	136
Rh/Y-zeolite, for propionitrile formation, <i>a</i>	92	Ru-Ir, (η <sup>4</sup> -C <sub>8</sub> H <sub>12</sub> )IrRu(μ-H)(PPh <sub>3</sub> ) <sub>2</sub> (η <sup>5</sup> -C <sub>2</sub> B <sub>9</sub> H <sub>11</sub> ), preparation, <i>a</i>	89
<b>Rhodium Alloys</b> , Rh-B/, Rh-Zn-B/SiO <sub>2</sub> , for hydroformylation of styrenes, <i>a</i>	140	<b>Coatings</b> , <i>a</i>	138
<b>Rhodium Complexes</b> , chiral, Rh, <i>a</i>	44	Nafion/glassy C disks + Os(bpy) <sub>3</sub> <sup>2+</sup> , electron transfer rates, <i>a</i>	90
(((cod)Rh(μ-Cl)) <sub>2</sub> ) + mono & bidentate ligands, for CO <sub>2</sub> hydrogenation, <i>a</i>	93	Pd, electroplated, on Ni-Co chip carriers, <i>a</i>	138
η <sup>5</sup> -CpRh(cod)/PS, for cyclopentanone synthesis, <i>a</i>	140	<b>Cobalt</b> , Co/Pt multilayers, coercivity, <i>a</i>	135
Rh <sub>2</sub> (OAc) <sub>4</sub> , Rh <sub>2</sub> (cap) <sub>4</sub> , Rh <sub>2</sub> (tfa) <sub>4</sub> , for α-diazo carbonyls reactions, <i>a</i>	93	CoPt single crystals, magnetism, <i>a</i>	38
Rh carbonyl/phenyl-sulphonato-alkylphosphine, for higher molecular olefin hydroformylation, <i>a</i>	44	Co-rich CoPt thin films, magnetic measurements, <i>a</i>	194
Rh phosphine PNS/PC, for hydroformylation, <i>a</i>	93	<b>Cold Start</b> , automotive, SAE conference	110
Rh(acac)(CO) <sub>2</sub> +phosphine ligands, for unsaturated alcohol hydroformylation, <i>a</i>	141	<b>Colloids</b> , Pd particles, ultrafine, preparation, <i>a</i>	142
RhCl <sub>3</sub> + poly- <i>N,N</i> -diallyl- <i>N,N</i> ,dimethylammonium chloride, for hexene hydroformylation, <i>a</i>	44	Pd(II) dispersion, electroless Ni deposition, <i>a</i>	138
RhCl <sub>3</sub> ·4H <sub>2</sub> O-(C <sub>8</sub> H <sub>17</sub> ) <sub>3</sub> N/Nd <sub>2</sub> O <sub>3</sub> , for hydrogenation	8		

	Page		Page
<b>Colloids (contd.)</b>		<b>Electrodes (contd.)</b>	
Pt, shape control	168	Pd, sheet, for styrene hydrogenation, <i>a</i>	137
for water photolysis, <i>a</i>	190	Pd-doped/C strips, printed, for HCHO sensing, <i>a</i>	42
Sb-doped SnO <sub>2</sub> , with Ru(bpy) <sub>3</sub> <sup>2+</sup> , luminescence, <i>a</i>	190	Pd/C paste, Pd immobilisation studies, <i>a</i>	137
<b>Composites, Pd/Ni wire, interdiffusion, <i>a</i></b>	88	platinised Pt, for enzyme detection, <i>a</i>	191
Pd/Ta, for H <sub>2</sub> separation, <i>a</i>	139	Pt(100), Pt(110), for HCHO electro-oxidation,	
thin Pd films/ceramic membranes, preparation, <i>a</i>	142	in perchloric acid, <i>a</i>	137
<b>Conferences, 2nd European Congress on Catalysis,</b>		Pt(111), HCHO surface patterns, <i>a</i>	40
Maastricht, September, 1995	19	hydride surface patterns, <i>a</i>	40
11th World Hydrogen Energy Conference,		Pt, bottom electrodes, on high dielectric SiO <sub>2</sub> /Si	
Stuttgart, 23-28 June 1996	175	thin films, for (Ba, Sr)TiO <sub>3</sub> films, <i>a</i>	45
Hydrogen Treatment of Materials,		for humidity sensors, <i>a</i>	138
Donetsk, September, 1995	36	polarised + PPy, NH <sub>3</sub> detection in biosensors, <i>a</i>	138
Metal-Hydrogen Systems,		polymer modified, for H <sub>2</sub> O <sub>2</sub> detection,	
Dayton, 10-11 June, 1996	180	with ascorbic acid, <i>a</i>	191
SAE, Spring, Detroit, 24-27 February 1996	110	Pt hydrous oxide film formation, <i>a</i>	39
6th Int. Conf. on the Chemistry of the Platinum		for D-sorbitol electro-oxidation, <i>a</i>	40
Group Metals, York, 22-26 July 1996	71, 161	to detect choline, acetylcholine	69
<b>Copolymerisation, ethylene and alkyl acrylates,</b>		Pt (nanosized)/graphite (porous),	
by Pd(II) catalyst	78	for O reduction reaction, <i>a</i>	89, 90
<b>Copper, plated onto Pd/carbon spheres, <i>a</i></b>	191	Pt and Pt-Ru, in DMFC	150
<b>Corrosion Protection, <i>a</i></b>	94	Pt single crystals, H deposition on, <i>a</i>	40
in the chemical and oil industries	54	NO adsorption, <i>a</i>	39
<b>Creep, in pure Pt, high temperature, <i>a</i></b>	189	Pt (well calibrated)/graphite (porous),	
<b>Cross-Coupling Reactions, of captodative olefins, <i>a</i></b>	193	for O reduction reaction, <i>a</i>	89
<b>Crotonaldehyde, hydrogenation,</b>		Pt-Ru/, Pt-Sn/, Pt-Ru-Sn/conducting polymer matrix,	
on Pt-Sn/C, Sn effect, <i>a</i>	139	poisoning for MeOH electro-oxidation, <i>a</i>	137
<b>Cyclisations, Heck-type, with Pd(OAc)<sub>2</sub>/TPPTS,</b>		Pt/C in PEFC, <i>a</i>	194
PdCl <sub>2</sub> /TPPTS, <i>a</i>	140	Pt/C/vulcan C, in fuel cells, <i>a</i>	141
<b>Cyclopentanone, synthesis, on η<sup>5</sup>-CpRh(cod)/PS, <i>a</i></b>	140	Pt/graphite, for MeOH electro-oxidation, <i>a</i>	89
<b>Cyclopropanation, ethyl diazoacetate with olefins, <i>a</i></b>	44	Pt/graphite/polymerised LaPO <sub>4</sub> , HER, <i>a</i>	137
		Pt/spherical microelectrodes, for CV	
<b>Dehydrocyclisation, <i>n</i>-hexane, <i>a</i></b>	192	and stripping analysis, <i>a</i>	42
<b>Dendralene, production, <i>a</i></b>	93	Pt/(Ta <sub>2</sub> O <sub>5</sub> +Pt)/Ti, fabrication, <i>a</i>	137
<b>Detectors see Sensors</b>		Pt/YSZ, in SOFC, <i>a</i>	194
<b>α-Diazo Carbonyls, reactions via Rh<sub>2</sub>(OAc)<sub>4</sub>,</b>		Pt/ZrO <sub>2</sub> , in O <sub>2</sub> automotive sensors, <i>a</i>	42
Rh <sub>2</sub> (cap) <sub>4</sub> , Rh <sub>2</sub> (tfa) <sub>4</sub> , <i>a</i>	93	PZT/Pt/SiO <sub>2</sub> /Si, (Ba, Sr)TiO <sub>3</sub> /Pt/SiO <sub>2</sub> /Si, thin films, <i>a</i>	90
<b>Dibenzothiophene, hydrodesulphurisation, <i>a</i></b>	139	Ru(NH <sub>3</sub> ) <sub>6</sub> <sup>3+</sup> and Os(bpy) <sub>3</sub> <sup>3+</sup> /Nafion,	
<b>Diesel, reactions with catalysts, <i>a</i></b>	92	for Ru oxidation to Ru(VII) by Os, <i>a</i>	137
<b>Diesel Fuel, hydrodesulphurisation, <i>a</i></b>	139	RuO <sub>2</sub> -SnO <sub>2</sub> coated Ti, morphology, <i>a</i>	90
<b>Dihydroxylation, alkenes, by RuCl<sub>3</sub>·(H<sub>2</sub>O)<sub>3</sub>, <i>a</i></b>	194	RuO <sub>2</sub> -TiO <sub>2</sub> film, SIMS study, <i>a</i>	138
<b>Diketones, β-Pt, <i>a</i></b>	189	RuO <sub>2</sub> /TiN, for (Ba, Sr)TiO <sub>3</sub> thin film capacitors, <i>a</i>	45
<b>Diodes, Pt/PZT/<i>n</i>-SrTiO<sub>3</sub> ferroelectric memory, <i>a</i></b>	142	RuO-RuCN/C for aliphatic,	
<b>syn-Diols, preparation, <i>a</i></b>	194	furanic aldehyde detection, <i>a</i>	42
<b>Diphenylacetylene, for indole formation, <i>a</i></b>	193	RuS <sub>2</sub> colloids/TiO <sub>2</sub> , photoelectrochemistry, <i>a</i>	41
<b>Diyne, synthesis, via Pd/TPPTS, <i>a</i></b>	44	<b>Electroless Plating, Pd/Vycor glass, bath conditions, <i>a</i></b>	42
<b>DNA, photochemical reactions with Ru(II), Rh(III)</b>		<b>Emission Control, at SAE conference</b>	110
polypyridyls, porphyrins, uranyl ions, <i>a</i>	91	<b>Enthalpies, of pgms</b>	62
Pt complexes in binding studies	127	<b>Entropies, of pgms</b>	62
<b>Electrical Contacts, ohmic,</b>		<b>Enzyme, electrodes, sensors, <i>a</i></b>	191
Au/Ge/Pd to <i>n</i> -type GaAs, <i>a</i>	194	<b>Ethane, ammoxidation for HCN synthesis, <i>a</i></b>	192
Pd/Al- <i>n</i> -type GaN, <i>a</i>	194	<b>Ethene, for direct alcohol formation, <i>a</i></b>	141
PdGe-based/ <i>n</i> -GaAs, thermal stability, <i>a</i>	45	for propionitrile formation, <i>a</i>	92
Ti/Pt/Au on Be implanted GaAsSb, HIGFET, <i>a</i>	94	<b>Ethylene, combustion on IrO<sub>2</sub>, NEMCA effect, <i>a</i></b>	40
<b>Electrocatalysts, for fuel cell</b>	94, 141, 142, 150, 194	for copolymerisation with CO, <i>a</i>	93
<b>Electrochemistry, <i>a</i></b>	39-40, 89-90, 137-138, 190	polymerisation, by Pd catalyst	78
<b>Electrodeposition, <i>a</i></b>	42, 138, 191	<b>Ethynyltrimethylsilane, polymerisation,</b>	
Cr, <i>a</i>	42	by Pd bis(acetylilide)complex, <i>a</i>	141
electroless metal plating, Pd on carbon spherules, <i>a</i>	191	<b>Extraction, noble metals from anode muds, in Russia</b>	118
Pd/Vycor glass, <i>a</i>	42	<b>Fluorescence, [PtBrMe<sub>2</sub>(4-CH<sub>2</sub>-7-MeO-coumarin)(phen)], <i>a</i></b>	39
<b>Electrodes, anodes, Pt/Ti for Cr electroplating, <i>a</i></b>	42	<b>Formaldehyde, detector, <i>a</i></b>	42
Pt, in PEFC, <i>a</i>	142	electro-oxidation, on Pt(100), Pt(110) electrodes,	
RuO <sub>2</sub> /TiO <sub>2</sub> , for sea water electrolysis, <i>a</i>	137	in perchloric acid, <i>a</i>	137
catalysts, Pt-Co-Cr/C black,		<b>Formic Acid, decomposition by</b>	
for O reduction reaction in PAFC, <i>a</i>	94	[Pt <sub>4</sub> (μ-CH <sub>3</sub> COO) <sub>8</sub> ]/SiO <sub>2</sub> , <i>a</i>	92
Pt/PAN, for O <sub>2</sub> reduction, <i>a</i>	94	production, via Rh complex catalyst, <i>a</i>	93
coated Ru-Pt oxide, for O <sub>2</sub> evolution and		<b>Fuel Cells</b>	44-45, 94, 141-142, 150, 194
hypochlorite production, <i>a</i>	90	at 11th World Hydrogen Energy Conference	175
enzyme, with Ir/C for biosensing glucose, <i>a</i>	191	book review	159
Ir-dispersed C-paste transducer biosensor, <i>a</i>	191	DMFC	150
IrO <sub>2</sub> coated Ti electrodes, O reduction kinetics, <i>a</i>	190	PAFC, Pt on electrodes, <i>a</i>	44
Ni-40Nb-pgm, for Zn extraction	18	PEFC, <i>a</i>	142
[Os(bipy) <sub>2</sub> (PVP) <sub>10</sub> Cl]Cl-modified, charge transport, <i>a</i>	190	gas diffusion electrodes, <i>a</i>	194
[Os(bpy)]/Nafion coatings, mass transport study, <i>a</i>	40	SOFC, <i>a</i>	45, 194

	<i>Page</i>		<i>Page</i>
<b>Fullerenes</b> , [C <sub>70</sub> (Pt(PPh <sub>3</sub> ) <sub>2</sub> ) <sub>4</sub> ], characterisation, <i>a</i>	189	<b>Hydrogenation</b> ( <i>contd.</i> )	
pgm complexes	23	propionaldehyde, <i>a</i>	194
Ru <sub>3</sub> -C <sub>60</sub> -Cs <sup>+</sup> /SiO <sub>2</sub> catalyst, for NH <sub>3</sub> synthesis, <i>a</i>	43	styrene, with Pd sheet electrode, for ethylbenzene, <i>a</i>	137
<b>Furanic Aldehyde</b> , detector, <i>a</i>	42	<b>Hydrogenolysis</b> , of chiral allylic and	
		propargylic cyclic carbonates, <i>a</i>	141
<b>Gallium Arsenide</b> , Ru, S passivation, <i>a</i>	88	<b>Hydroxylation</b> , alkanes, with O <sub>2</sub> and	
<b>Gauzes</b> , single Pt-Rh, for rapid alkane conversions	125	Ru polyoxometalate, <i>a</i>	44
<b>Glucose</b> , sensors, <i>a</i>	91, 191	<b>Hypochlorite</b> , production, on Ru-Pt oxide electrodes, <i>a</i>	90
<b>Graphite</b> , in Pt(IV)-graphite compounds, <i>a</i>	40		
		<b>Indoles</b> , formation from azobenzene and	
<b>Hexanal</b> , photocatalytic hydrogenation,		diphenylacetylene, <i>a</i>	193
with H <sub>2</sub> by RhCl(PMe <sub>3</sub> ) <sub>2</sub> CO, <i>a</i>	190	<b>Iridium</b> , apparatus, for XRF fusion technology	87
<b>n-Hexane</b> , conversions on Pt-Dy/Al <sub>2</sub> O <sub>3</sub> , <i>a</i>	192	compounds, NaSr <sub>3</sub> IrO <sub>6</sub> , single crystals, <i>a</i>	136
<b>Hexene-1</b> , hydroformylation, <i>a</i>	44	Sr <sub>3</sub> CuPt <sub>1-x</sub> Ir <sub>x</sub> O <sub>6</sub> , random quantum spin	
<b>History</b> , discovery of ruthenium	181	chain paramagnetism	63
ruthenium, and recent research	128	SrCaIrO, SrBaIrO, properties, <i>a</i>	38
<b>Humidity</b> , sensor, <i>a</i>	138	contact electroresistance,	
<b>Hydration</b> , of nitriles to amides	169	for oxide film defect evaluation, <i>a</i>	91
<b>Hydrocarbonylation</b> , alkenes, from alcohols, <i>a</i>	141	<b>Iridium Alloys</b> , Ir-Nb, Ir-Ti, refractory superalloys, <i>a</i>	189
in platinum metals metallurgy	118	<b>Iridium Complexes</b>	
<b>Hydrodesulphurisation</b> , benzothiothiophene and diesel, <i>a</i>	139	(η <sup>4</sup> -C <sub>8</sub> H <sub>12</sub> )IrRu(μ-H)(PPh <sub>3</sub> ) <sub>2</sub> (η <sup>5</sup> -C <sub>2</sub> B <sub>9</sub> H <sub>11</sub> ) cluster, <i>a</i>	89
<b>Hydroformylation</b> , alcohols, unsaturated,		[Ir(COD)Cl] <sub>2</sub> + α-diaimines, <i>a</i>	39
by Rh(acac)(CO) <sub>2</sub> +phosphine ligands, <i>a</i>	141	<b>Iridium Oxide</b> , catalyst film, for ethylene combustion,	
asymmetric, styrene, <i>a</i>	43, 44	NEMCA effect, <i>a</i>	40
by CpRh(cod)/PS, <i>a</i>	140	compounds, electrochemistry during O reactions, <i>a</i>	40
hex-1-ene to aldehydes, alcohols, by [RhH(PEt <sub>3</sub> ) <sub>3</sub> ], <i>a</i>	141	<b>Iridium Silicides</b> , Si:Ir deposition ratio effects	
hexene-1, <i>a</i>	44	on IrSi/Si(100), <i>a</i>	135
higher molecular olefins,		<b>Iron</b> , [Fe(II),Ru(III)] formation, <i>a</i>	94
by H <sub>2</sub> O + MeOH soluble Rh catalysts, <i>a</i>	44		
liquid-phase, styrene, substituted, <i>a</i>	140	<b>Johnson Matthey</b> , "Platinum 1996"	127
methylacrylate to α-aldehyde,		relationship with Rustenburg Platinum Mines	2
via water soluble Rh phosphine PNS/PC, <i>a</i>	93	<b>Junctions</b> , Pt silicided p <sup>n</sup> , Pt/Si films/Si, <i>a</i>	45
<b>Hydrogen</b> , changing mechanical properties and			
structure of Pd-Mn	37	<b>Ketone</b> , production, from olefin C=C bond fission,	
deposition on single crystal Pt, Pt(100), Pt(110),	40	in presence of K osmate, <i>a</i>	90
and Pt(111) surfaces, <i>a</i>		<b>Klaus, Karl Karlovitch</b> , discoverer of ruthenium	181
evolution reaction, on electrodes of			
Pt/graphite/polymerised LaPO <sub>4</sub> , <i>a</i>	137	<b>Lactones</b> , from 2-alkynylbenzaldehyde carbonylation, <i>a</i>	193
forced diffusion in Pd, <i>a</i>	189	<b>Lanthanum</b> , additions to 3-way automotive catalysts, <i>a</i>	192
interaction with Pd, Donetsk conf. report	36	<b>Lean-Burn Technology</b> , at SAE conference	110
permeation in Pd-Ag membranes, <i>a</i>	38	<b>Linoleic Acid</b> , photohydrogenation by Pt/TiO <sub>2</sub> , <i>a</i>	41
photoproduction, on Pt/CdS, Pt/CdS/ZnS, <i>a</i>	90	<b>LPG</b> , sensor, <i>a</i>	191
photoproduction from water,		<b>Luminescence</b> ,	
by Pt/TiO <sub>2</sub> +Zn porphyrin, <i>a</i>	41	in [Ru(bpy) <sub>2</sub> (bpz)] <sup>2+</sup> , in [Zn(bpy) <sub>3</sub> ](ClO <sub>4</sub> ) <sub>2</sub> , <i>a</i>	138
by Rh/perfluorinated poly( <i>p</i> -phenylene), <i>a</i>	41	of Ru(bpy) <sub>3</sub> <sup>2+</sup> on colloidal Sb-doped SnO <sub>2</sub> , <i>a</i>	190
by water photolysis, <i>a</i>	190	[Ru(bpy-d <sub>3</sub> ) <sub>2</sub> (bpz)] <sup>2+</sup> , in [Zn(bpy) <sub>3</sub> ](ClO <sub>4</sub> ) <sub>2</sub> , <i>a</i>	138
discussed at 11th World Hydrogen Energy Conf.	175	in Ru(bpy)(CN), Ru(DMbp)(CN), <i>a</i>	41
in a PEFC, <i>a</i>	142	<b>L-Lysine</b> , photocatalysed to pipecolic acid, <i>a</i>	41
reaction with CO over Fe + Pt/SiO <sub>2</sub> , <i>a</i>	43		
sensors, <i>a</i>	42, 191	<b>Magnetism</b> , in Co-rich CoPt and CoPtP thin films, <i>a</i>	194
separation, <i>a</i>	139	coercivity, Co/Pt multilayers, <i>a</i>	135
solubility in Pd-Pt, <i>a</i>	88	in CoPt single crystals, under compression, <i>a</i>	38
storage, for fuel cell use	177	electronic and nuclear in Pt	112
<b>Hydrogen Cyanide</b> , synthesis, <i>a</i>	192	in FeRhPd alloys, and crystalline properties, <i>a</i>	88
<b>Hydrogen Peroxide</b> , detection, <i>a</i>	191	in LaNdRhSi system, <i>a</i>	38
<b>Hydrogenation</b> , 1,3-butadiene, over Co-Pd/Al <sub>2</sub> O <sub>3</sub> , <i>a</i>	43	in MnPt thin films, <i>a</i>	135
acrylic acid, on Pd/PA-Nd <sup>3+</sup> , <i>a</i>	192	random quantum spin chain paramagnetism	
alkenes, on Pt/Na-BEA, <i>a</i>	139	in Sr <sub>3</sub> CuPt <sub>1-x</sub> Ir <sub>x</sub> O <sub>6</sub>	63
asymmetric transfer, of ketones, in propan-2-ol,		in SrCaIrO, SrBaIrO, structures, <i>a</i>	38
via Ru(II) complex, <i>a</i>	94	in UIr <sub>2</sub> Si <sub>2</sub> , and physical properties, <i>a</i>	88
cinnamaldehyde, over Rh, Ru, Pd complexes, <i>a</i>	192	in U(PdFe) <sub>2</sub> Ge <sub>3</sub> , <i>a</i>	88
citral, on Ru-Me/Al <sub>2</sub> O <sub>3</sub> , Me = Ge, Sn, Pb, <i>a</i>	140	<b>Manganese</b> , MnPt thin films, magnetic properties, <i>a</i>	135
CO <sub>2</sub> to EtOH, over Rh/SiO <sub>2</sub> +Li salts, <i>a</i>	193	<b>Medical</b> , carboplatin, photostudies, <i>a</i>	138
CO <sub>2</sub> to HCOOH, <i>a</i>	93	corrosion of implants, <i>a</i>	94
CO <sub>2</sub> by Pt/Co/Al <sub>2</sub> O <sub>3</sub> , <i>a</i>	43	<b>Medical Uses</b> , <i>a</i>	45, 165
on Mn-Ru/Al <sub>2</sub> O <sub>3</sub> catalysts, <i>a</i>	140	<b>Melting Points</b> , of pgms	62
crotonaldehyde, on Pt-Sn/C, Sn effect, <i>a</i>	139	<b>Membranes</b> , Pd-Ag films, preparation, <i>a</i>	38
hexanal, with H <sub>2</sub> in RhCl(PMe <sub>3</sub> ) <sub>2</sub> CO, <i>a</i>	190	Pd/α-Al <sub>2</sub> O <sub>3</sub> , tubes, for H <sub>2</sub> removal, <i>a</i>	192
linoleic acid, on Pt/TiO <sub>2</sub> , <i>a</i>	41	Pd/ceramic composite, preparation, <i>a</i>	142
liquid-phase, by pgm complex catalysts	8	Pd/Ta composites, for H <sub>2</sub> separation, <i>a</i>	139
of C <sub>6</sub> olefinic hydrocarbons, on Pd/AFC, <i>a</i>	140	Pt nanochannel replicas	25
naphthalene, with Pt-Bx-Al <sub>2</sub> O <sub>3</sub> , <i>a</i>	191	<b>Memory Diode</b> , Pt/PZT/n-SrTiO <sub>3</sub> , fabrication, <i>a</i>	142
on Pt/AAPx catalyst, <i>a</i>	42	<b>Methane</b> , ammoxidation, to HCN on	
organic substrates, by Rh(PPh <sub>3</sub> ) <sub>3</sub> Cl-isocyanide		Pt/Al <sub>2</sub> O <sub>3</sub> , Pt-Rh, <i>a</i>	192
polymer bound, <i>a</i>	193	conversion to syngas over Pt electrode, <i>a</i>	194

	Page		Page
<b>Methylacrylate</b> , hydroformylation to $\alpha$ -aldehyde, via water soluble Rh phosphine PNS/PC, <i>a</i>	93	<b>Palladium Alloys</b> , Ag-Pd, Cu-Pd, nanoscale, <i>a</i>	38
<b>Methylcyclopentane</b> , conversions, <i>a</i>	192	Cu-Pd, properties and structures, <i>a</i>	189
<b>Mining</b> , platinum, in South Africa	2	Pd-Ag, film membranes, preparation, <i>a</i>	38
<b>Mirrors</b> , X-ray multilayer, Pt/C, <i>a</i>	88	Pd-Cu, CO, NO surface reactions, <i>a</i>	38
<b>Modifiers</b> , Ba, for 3-way catalysts, <i>a</i>	92, 192	Pd-Mn, H changes mechanical properties, structure	37
<b>Molecular Wires</b>	72	Pd-Pt, thick foil, H solubility, <i>a</i>	88
Ru and Os polypyridine complexes	26	<b>Palladium Complexes</b> , acetylacetonato derivatives	
<b>Molybdenum</b> , effect on corrosion resistance of V-free Ti medical implants, <i>a</i>	94	of palladated $\alpha$ -arylalkylamines, synthesis, <i>a</i>	136
<b>Nanocomposites</b> , Pt/polymer preparation	77	$(\eta^2\text{-C}_{60})\text{Pd}(\text{CNR})_2$ , <i>a</i>	189
<b>Nanoparticles</b> , Pt in recast ionomer, for O reduction, <i>a</i>	89	$(\eta^2\text{-C}_{60})\text{Pd}(\text{PPh}_3)_2$ , in benzene, laser excitation spectra, <i>a</i>	190
<b>Nanotechnology</b> , molecular wires	26	palladacycle, from $\text{Pd}(\text{PPh}_3)_2\text{Cl}_2$ + dimethylhydrazone, <i>a</i>	39
nanometre Pt/MCM-41	35	Pd macrocycles, with pendant-arm 14-,16-,18-members, <i>a</i>	189
Pt nanochannel replica membranes	25	$\text{Pd}(\text{NH}_3)_4^{2+}$ ions, on C spheres, for Ni, Cu plating, <i>a</i>	191
<b>Naphthalene</b> , hydrogenation, on Pt-Bx-Al <sub>2</sub> O <sub>3</sub> catalyst, <i>a</i>	191	polynuclear hydroxo Pd <sup>II</sup> (PHC) complexes, <i>a</i>	39
on Pt/AAPx catalyst, <i>a</i>	42	<b>Permeation</b> , H, in Pd	40, 175, 180
<b>NEMCA</b> , effect, <i>a</i>	40	H in Pd-Ag membranes, <i>a</i>	38
<b>New Patents</b> , <i>a</i>	46-52, 95-100, 143-148, 195-198	<b>Phase Diagrams</b> , RuO <sub>2</sub> -Bi <sub>2</sub> O <sub>3</sub> -ZnO, RuO <sub>2</sub> -Al <sub>2</sub> O <sub>3</sub> -ZnO, <i>a</i>	88
<b>Nickel</b> , electroless deposition, via Pd(II) colloid, <i>a</i>	138	<b>Photocatalysts</b> , Pt/CdS, L-lysine into pipecolic acid, <i>a</i>	41
plated onto Pd/carbonaceous spherules, <i>a</i>	191	Pt/TiO <sub>2</sub> , for H <sub>2</sub> production from water, <i>a</i>	41
<b>Nitric Acid</b> , plants	80	for linoleic acid hydrogenation, <i>a</i>	41
<b>Nitriles</b> , hydration to amides	169	PtS/CdS/Pt, IrS/CdS/Ir, <i>a</i>	41
<b>Nitriloacetic Acid</b> , electron donor, for H <sub>2</sub> photoproduction, <i>a</i>	190	Rh colloid/perfluorinated poly( <i>p</i> -phenylene), <i>a</i>	41
<b>Nitrogen Oxides</b> , NO,		<b>Photoconversion</b> , <i>a</i>	41, 90, 91, 138, 190
adlayers on Pt single crystal electrodes, <i>a</i>	39	<b>Photolysis</b> , for H <sub>2</sub> photoproduction, <i>a</i>	190
and CO adsorption on Rh/lanthana, Pt/lanthana, <i>a</i>	91	laser, of $(\eta^2\text{-C}_{60})\text{Pd}(\text{PPh}_3)_2$ , spectra, <i>a</i>	190
decomposition using C <sub>2</sub> H <sub>6</sub> , H <sub>2</sub> , over Pt/Al <sub>2</sub> O <sub>3</sub> , <i>a</i>	91	<b>Photoreactions</b> , with DNA, induced by Ru(II), Rh(III), polypyridyls, porphyrins, uranyl ions, <i>a</i>	91
reactions with CO, on La <sub>0.8</sub> Sr <sub>0.2</sub> Al <sub>0.8</sub> Cu <sub>0.1</sub> Ru <sub>0.1</sub> O <sub>3</sub> , <i>a</i>	193	<b>Plating</b> , Ni onto Pd/carbonaceous spherules, <i>a</i>	191
reactions with PdCu, <i>a</i>	38	<b>"Platinum 1996"</b>	127
<b>NOx</b> , reductions, by 3-way catalysts, <i>a</i>	92	<b>Platinum</b> , additions, to YBaCuO superconductors, effects on solidification, <i>a</i>	135
reductions over Pd modified 3-way catalysts, <i>a</i>	192	on CdS and CdS/ZnS, for H <sub>2</sub> photoproduction, <i>a</i>	90
sensor, with Pt gate MOSFET, <i>a</i>	91	Co/Pt multilayers, coercivity, <i>a</i>	135
<b>Olefins</b> , higher molecular, hydroformylation, <i>a</i>	44	colloids	168, 190
$\alpha$ -Olefins, polymerisation, by Pd catalyst	78	compounds, Pt chloride-graphite intercalation, <i>a</i>	40
<b>Optical Switching</b> , in Pd/Y and Pd/La films	102	Sr <sub>3</sub> CuPt <sub>1-x</sub> Ir <sub>x</sub> O <sub>6</sub> , random quantum spin chain paramagnetism	63
<b>Osmium</b> , compounds, LaRu <sub>4-x</sub> Os <sub>2</sub> P <sub>12</sub> , superconductivity, electrical conductivity, <i>a</i>	135	Tl <sub>2</sub> Pt <sub>2</sub> S <sub>6</sub> , Tl <sub>2</sub> Pt <sub>4</sub> Te <sub>6</sub> , Tl <sub>2</sub> Pt <sub>4</sub> Se <sub>6</sub> , preparation, <i>a</i>	89
<b>Osmium Complexes</b> , chirals, Os(CHIRAGEN[7])	102	contact electroresistance measurements, for oxide film defect evaluation, <i>a</i>	91
Os polypyridines, in molecular wires	26	in fuel cells, see <b>Fuel Cells</b>	
[Os(bipy) <sub>2</sub> (PVP) <sub>10</sub> Cl]Cl, thin films on electrodes, <i>a</i>	190	high temperature creep, <i>a</i>	189
[Os(bpy)] <sub>3</sub> <sup>2+</sup> , in Nafion coatings on glassy C disks, electron transfer rates, <i>a</i>	90	membranes	25
Os(bpy) <sub>3</sub> <sup>3+</sup> /Nafion, for oxidising Ru(NH <sub>3</sub> ) <sub>6</sub> <sup>3+</sup> , <i>a</i>	137	mining in South Africa	2
[Os(CO) <sub>6</sub> ][Sb <sub>2</sub> F <sub>11</sub> l <sub>2</sub> ], <i>a</i>	39	Pt <sub>309</sub> clusters, with ligands, inner metallic core, <i>a</i>	136
sandwich indenyl, $(\eta^2\text{-C}_9\text{H}_7)\text{Os-}\eta^5\text{-C}_9\text{H}_7$ , redox behaviour at a Pt electrode, <i>a</i>	90	Pt hydrous oxide films, from potential cycling, <i>a</i>	39
used in molecular wire development	26, 72	PtCo single crystals, magnetism, <i>a</i>	38
<b>3-Oxabicyclo[4.1.0]hept-4-enes</b> , synthesis by PtCl <sub>4</sub> , <i>a</i>	43	recovery in nitric acid plants	80
<b>Oxidation</b> , alkanes, by RuF <sub>10</sub> Pc in NaX zeolite catalyst	37	in superconductivity	64
CO, on Pt/CeO <sub>2</sub> , kinetic modelling, <i>a</i>	139	<b>Platinum Alloys</b> , Pt <sub>2</sub> Sn, single crystal surfaces for CO, CO:H <sub>2</sub> electro-oxidation, <i>a</i>	90
electro, MeOH, on Pt/graphite, <i>a</i>	89	<b>Platinum Complexes</b> , [C <sub>70</sub> {Pt(PPh <sub>3</sub> ) <sub>2</sub> }] <sub>4</sub> , isolation, characterisation, <i>a</i>	189
partial, alkanes, Pt-Rh gauze	125	chiral complexes	102
<b>Oxygen</b> , dissolved, sensor, <i>a</i>	191	$(\eta^2\text{-C}_{60})\text{Pt}(\text{CNR})_2$ , <i>a</i>	189
evolution on coated Ru-Pt oxide electrodes, <i>a</i>	90	[Pt <sub>2</sub> ( $\mu$ -Cl) <sub>2</sub> ]{(COR) <sub>2</sub> H <sub>2</sub> }, R=Me, Et, <i>a</i>	189
reduction, in alkali, on IrO <sub>2</sub> /Ti electrodes, <i>a</i>	190	[PtBrMe <sub>2</sub> (4-CH <sub>2</sub> -7-MeO-coumarin)(phen)], fluorescence, <i>a</i>	39
<b>Palladium</b> , coatings on tubular reactors, <i>a</i>	192	[Pt <sup>II</sup> (dpp-C,N,N')(MeCN)] <sup>+</sup> , DNA binding studies	127
compounds, Pd in VOHPO <sub>4</sub> ·0.5H <sub>2</sub> O, <i>a</i>	89	<i>cis</i> -Pt(LH <sub>2</sub> )Cl <sub>2</sub> , H <sub>2</sub> L[PtCl <sub>4</sub> ], with deprotonated amide ligands, <i>a</i>	39
Tl <sub>2</sub> Pd <sub>4</sub> Se <sub>6</sub> , preparation, <i>a</i>	89	<i>trans</i> -Pt(II) acetylide, photophysical properties, <i>a</i>	91
corrosion effects on V-free Ti alloys for implants, <i>a</i>	94	<b>Platinum Silicide</b> , in junctions, <i>a</i>	45
electrocoatings for Ni-Co chip carriers, <i>a</i>	138	<b>Platinum-Iron</b> , ultralow temperature physics	112
forced H diffusion, <i>a</i>	189	<b>Poisoning</b> , Pt-based electrodes,	
H diffusion through sheet electrode, <i>a</i>	137	in MeOH electro-oxidation, <i>a</i>	137
H permeation, <i>a</i>	40	of Pt-Sn/Al <sub>2</sub> O <sub>3</sub> by thiophene, <i>a</i>	192
nanocrystalline, specific heat, <i>a</i>	38	S in fuel cells, <i>a</i>	141
Pd-H, anomaly at 50K	180	<b>Polymerisation</b> , CO + ethylene, via Pd complex, <i>a</i>	93
Donetsk conf.	36	ethynyltrimethylsilane, by Pd bis(acetylide), <i>a</i>	141
system, ACS symposium	180	<b>Polymers</b> , high molecular weight, production, by Pd catalyst system	78
thin films on Y and La, optical switching effect	102	with Os complex as electrode coating, <i>a</i>	190
ultrafine particles, preparation using ultrasound, <i>a</i>	142	polyketone, production via (dppp)Pd(OAc) <sub>2</sub> or (dppp)Pd[C(O) <sup>t</sup> Bu]Cl + <i>tert</i> -butyl alumoxane, <i>a</i>	93
ultrathin films, growth on Al(001), <i>a</i>	135		

	Page		Page
<b>Printing</b> , fine electronic circuitry	116	<b>Sensors (contd.)</b>	
<b>Promoters</b> , CeO <sub>2</sub> , for 3-way catalyst, <i>a</i>	92	NOx, with Pt gate MOSFET, <i>a</i>	91
Li salts for Rh/SiO <sub>2</sub> , CO, hydrogenation to EtOH, <i>a</i>	193	O, dissolved, +fluorescent quenching of Ru complex, <i>a</i>	191
polymer-immobilised Nd ions, for ultrafine Pd, <i>a</i>	192	<b>Single Crystals</b> , CoPt, magnetism in, <i>a</i>	38
<b>Propan-1-ol</b> , preparation, <i>a</i>	194	NaSr <sub>3</sub> IrO <sub>6</sub> , structure, <i>a</i>	136
<b>Propane</b> , aromatisation on Ga/SiO <sub>2</sub> ,		PdCu, CO, NO surface reactions, <i>a</i>	38
in Pd membrane reactor, <i>a</i>	192	Pt,Sn surfaces, for CO, CO:H <sub>2</sub> electro-oxidation, <i>a</i>	90
<b>Propionitrile</b> , formation, over Rh/Y-zeolite, <i>a</i>	92	Pt, hydride surface patterns, <i>a</i>	40
		Pt electrodes, NO adsorption, <i>a</i>	39
<b>Reactors</b> , tubular, Pd coated, for H removal, <i>a</i>	192	<b>Sol-Gel Process</b> , for O sensor production, <i>a</i>	191
<b>Recording Film</b> , ultra-high density, CoCrPt(Ta, B), <i>a</i>	45	for Pt-Sn/Al <sub>2</sub> O <sub>3</sub> catalysts, properties, <i>a</i>	192
<b>Reduction</b> , oxygen, on IrO <sub>2</sub> coated Ti electrodes, <i>a</i>	190	for Pt/TiO <sub>2</sub> catalyst preparation, characterisation, <i>a</i>	139
<b>Rhodium</b> , compounds, CeRh <sub>2</sub> Si <sub>2</sub> , superconductivity, <i>a</i>	135	<b>D-Sorbitol</b> , electro-oxidation on Pt, <i>a</i>	40
LaNdRhSi system, superconductivity, magnetism, <i>a</i>	38	<b>Specific Heats</b> , in nano-Pd, <i>a</i>	38
in superconductivity	64	of pgms	62
<b>Rhodium Complexes</b> , [c-BHP(C <sub>12</sub> ) <sub>2</sub> Rh <sup>III</sup> (L)(H <sub>2</sub> O)]Cl,		<b>Stearic Acid</b> , production, <i>a</i>	41
synthesis, characterisation, <i>a</i>	136	<b>Styrene</b> , asymmetric hydroformylation, <i>a</i>	43, 44
Rh <sup>I</sup> -Rh <sup>III</sup> with cyanide bridges, <i>a</i>	136	hydrogenation, with a Pd sheet electrode, <i>a</i>	137
<i>trans</i> -[Rh(CH=CH <sub>2</sub> )(CS)(PPr <sub>3</sub> ) <sub>2</sub> ], preparation, <i>a</i>	89	substituted, liquid-phase hydroformylation,	
<i>trans</i> -[Rh(R)(CO)(PPr <sub>3</sub> ) <sub>2</sub> ], preparation, <i>a</i>	89	on Rh-B, Rh-Zn-B/SiO <sub>2</sub> catalysts, <i>a</i>	140
<b>Rustenburg Platinum Mines</b>	2	<b>Sulphur</b> , catalyst tolerance, <i>a</i>	43
<b>Ruthenium</b> , compounds, CeRu <sub>2</sub> , superconductivity, <i>a</i>	89	effects on Pt electrocatalyst in fuel cell, <i>a</i>	141
CeRu <sub>4</sub> P <sub>12</sub> , superconductivity, electrical properties, <i>a</i>	135	S, catalyst resistance to, <i>a</i>	92
LaRu <sub>4</sub> Os <sub>2</sub> P <sub>12</sub> , superconductivity, <i>a</i>	135	<b>Superalloys</b> , Ir-Nb, Ir-Ti refractories, <i>a</i>	189
Ru(bpy)(CN), Ru(DMbp)(CN), photoproperties, <i>a</i>	41	<b>Superconductivity</b> , CeRu <sub>2</sub> , single crystal, <i>a</i>	89
Ru(NH <sub>3</sub> ) <sub>3</sub> <sup>3+</sup> , oxidation by Os(bpy) <sub>3</sub> <sup>3+</sup> /Nafion, <i>a</i>	137	flux pinning by Pt and Rh in YBaCuO	64
Ru[R <sub>2</sub> PCH <sub>2</sub> CH <sub>2</sub> PR <sub>2</sub> ] <sub>2</sub> H <sub>2</sub> , photoproducts,		in LaRu <sub>4</sub> P <sub>12</sub> , LaRu <sub>4</sub> OsP <sub>12</sub> , LaRu <sub>2</sub> Os <sub>2</sub> P <sub>12</sub> , <i>a</i>	135
after laser flash photolysis, <i>a</i>	41	in LaNdRhSi system, <i>a</i>	38
[Fe(II),Ru(III)] formation, <i>a</i>	94	<b>Superconductors</b> , CeRh <sub>2</sub> Si <sub>2</sub> , <i>a</i>	135
historical discovery	181	Sr <sub>2</sub> RuO <sub>4</sub> , quantum oscillations, <i>a</i>	136
historical perspective	128	UPd <sub>2</sub> Al <sub>3</sub> , crystal field transitions, <i>a</i>	135
<b>Ruthenium Alloys</b> , Ru enhanced Ti alloys	54	YBaCuO, effects of Pt additions to solidification, <i>a</i>	135
<b>Ruthenium Chloride</b> , to passivate GaAs surface, <i>a</i>	88	<b>Synthesis Gas</b> , preparation, <i>a</i>	194
<b>Ruthenium Complexes</b> , chiral, Ru(CHIRAGEN[6])Cl <sub>2</sub>	102		
( $\eta^5$ -C <sub>9</sub> H <sub>7</sub> ) <sub>2</sub> IrRu( $\mu$ -H)(PPh) <sub>2</sub> ( $\eta^5$ -C <sub>2</sub> B <sub>9</sub> H <sub>11</sub> ) cluster, <i>a</i>	89	<b>Temperature Measurement</b> , by Ir-Au proximity	
[Ru <sub>3</sub> (O <sub>2</sub> CCMe <sub>3</sub> )(NITPh)](BF <sub>4</sub> ) <sub>n</sub> , properties, <i>a</i>	136	effect thermometers, 20–100 mK, <i>a</i>	142
Ru <sub>3</sub> (CO) <sub>8</sub> (C <sub>9</sub> H <sub>7</sub> ), C-H bond activation		<b>Thermodynamic Properties</b> , Pt, Pd, Rh, Ir, Os, Ru	38, 62
on Ru-Ru fission, <i>a</i>	136	<b>Thick Films</b> , Pd-Pt, H solubility, <i>a</i>	88
Ru <sub>5</sub> ( $\mu_2$ -C <sub>2</sub> )( $\mu$ -SMe)( $\mu$ -PBu <sub>3</sub> )(CO) <sub>11</sub> ,		<b>Thin Films</b> , capacitors, <i>a</i>	45
preparation, characterisation, <i>a</i>	190	CoCrPt(Ta, B) as ultra-high density recording media, <i>a</i>	45
Ru polypyridines, in molecular wires	26	CoPtP alloys, magnetic values, <i>a</i>	194
Ru(bipy) <sub>3</sub> <sup>2+</sup> , in water photolysis system, <i>a</i>	190	high dielectric, Pt bottom electrodes on SiO <sub>2</sub> /Si,	
[Ru(bpy) <sub>2</sub> (bpz)] <sup>2+</sup> , [Ru(bpy) <sub>2</sub> (bpz)] <sup>2+</sup> , in		for (Ba, Sr)TiO <sub>3</sub> films, <i>a</i>	45
[Zn(bpy) <sub>3</sub> ](ClO <sub>4</sub> ) <sub>2</sub> , luminescence spectra, <i>a</i>	138	MnPt, magnetic properties, long range order, <i>a</i>	135
Ru(bpy) <sub>2</sub> (dcbpy) <sup>2+</sup> , electron transfer to SnO <sub>2</sub>		[Os(bipy) <sub>2</sub> (PVP) <sub>10</sub> Cl]Cl on electrodes,	
nanocrystallites, <i>a</i>	138	charge transport, <i>a</i>	190
Ru(bpy) <sub>2</sub> <sup>2+</sup> , on colloidal Sb-SnO <sub>2</sub> , luminescence, <i>a</i>	190	Pd, ultrathin, Al(110), growth, <i>a</i>	135
RuCl <sub>3</sub> ·3H <sub>2</sub> O in polycarbonate, interactions, <i>a</i>	93	Pd on ceramic composite membranes, <i>a</i>	142
[Ru(CO) <sub>6</sub> ][Sb <sub>2</sub> F <sub>11</sub> ] <sub>2</sub> , <i>a</i>	39	Pd/Y and Pd/La, optical switching	102
[Ru <sup>II</sup> tris(4,7-diphenyl-1,10-phenanthroline)]		PZT/Pt/SiO <sub>2</sub> /Si, (Ba, Sr)TiO <sub>3</sub> /Pt/SiO <sub>2</sub> /Si electrodes, <i>a</i>	90
in sol-gel SiO <sub>2</sub> , for O sensor, <i>a</i>	191	RuO <sub>2</sub> /YSZ, electrical properties, <i>a</i>	38
[Ru(NH <sub>3</sub> ) <sub>5</sub> Cl]Cl <sub>3</sub> , fac-[Ru(NH <sub>3</sub> ) <sub>3</sub> Cl <sub>3</sub> ], preparation, <i>a</i>	89	SnO <sub>2</sub> , Pd 2–5 nm, activated, gas sensing, <i>a</i>	42
sandwich indenyl, ( $\eta^5$ -C <sub>9</sub> H <sub>7</sub> )Ru- $\eta^5$ -C <sub>9</sub> H <sub>7</sub> ,		SnO <sub>2</sub> nanocrystalline,	
redox behaviour at a Pt electrode, <i>a</i>	90	electron transfer from Ru(bpy) <sub>2</sub> (dcbpy) <sup>2+</sup> , <i>a</i>	138
<i>trans</i> -[Ru(bpy) <sub>2</sub> (py <sup>+</sup> )(py <sup>+</sup> )] <sup>III</sup> ,		<b>Tin</b> , effects on Pt-Sn/C catalysts,	
chromophore-quencher complexes, <i>a</i>	190	for crotonaldehyde hydrogenation, <i>a</i>	139
used in molecular wire development	26, 72	<b>Tin Oxide</b> , nanocrystalline films,	
<b>Ruthenium Oxides</b> , RuO <sub>2</sub> grown on YSZ, <i>a</i>	38	electron transfer from Ru(bpy) <sub>2</sub> (dcbpy) <sup>2+</sup> , <i>a</i>	138
Sr <sub>2</sub> RuO <sub>4</sub> , superconductivity, quantum oscillations, <i>a</i>	136	SnO <sub>2</sub> , Pd thin films	42
		<b>Titanium Alloys</b> , medical implants, Mo, Pd effects, <i>a</i>	94
<b>Schottky Diodes</b> , Pt-InAlAs, Fe doped,		Ru enhanced	54
IVT characteristics, <i>a</i>	142	<b>Tunnel Junctions</b> , H <sub>2</sub> sensor, <i>a</i>	191
PtSi/p-Si, IR, bias and thickness dependence, <i>a</i>	94		
PtSi/Si(111), non-uniform barrier height effects, <i>a</i>	45	<b>Vapour Pressures</b> , of pgms	62
<b>Sensors</b> , ammonia, in biosensor, <i>a</i>	138	<b>Voltresistometry</b> , with Pt, Ir, for oxide film evaluation, <i>a</i>	91
by Pt-doped CdSnO <sub>3</sub> , <i>a</i>	191		
automotive, development of	110	<b>Water</b> , electro-oxidation by Ru/Pt	87
O <sub>2</sub> sensors, <i>a</i>	42	photodecomposition, <i>a</i>	41
enzyme, <i>a</i>	191	sea, electrolysis, via RuO <sub>2</sub> /TiO <sub>2</sub> , <i>a</i>	137
formaldehyde, <i>a</i>	42	<b>Wires</b> , composite Pd/Ni, interdiffusion, <i>a</i>	88
glucose, <i>a</i>	91, 191	molecular, development	26, 72
H <sub>2</sub> , <i>a</i>	42, 191		
H <sub>2</sub> O <sub>2</sub> , on polymer-modified Pt electrode, <i>a</i>	191	<b>m-Xylene</b> , total combustion over Pd/stainless steel, <i>a</i>	43
humidity, with Pt electrodes, <i>a</i>	138	<b>Zinc</b> , extraction with Nb-Ni-pgm electrode	18
LPG, <i>a</i>	191		